

G 18001031



18001031

Reg. No.....

Name.....

M.Sc. DEGREE (C.S.S.) EXAMINATION, MAY 2018

Fourth Semester

Faculty of Science

Branch—I (A)—Mathematics

MT 04 E 01—ANALYTIC NUMBER THEORY

(2012 Admission onwards)

Time : Three Hours

Maximum Weight : 30

Part A

*Answer any five questions.
Each question carries weight 1.*

1. Show that $\phi(p^\alpha) = p^\alpha - p^{\alpha-1}$ for p , prime and $\alpha \geq 1$.
2. Define Bell series of an arithmetical function. Also Find $\mu_p(x)$.
3. Show that $\sum_{n \leq x} \frac{1}{n} = \log x + c + o\left(\frac{1}{x}\right)$ if $x \geq 1$.
4. Show that $0 \leq \frac{4(x)}{x} - \frac{\mathcal{J}(x)}{(x)} \leq \frac{(\log x)^2}{\sqrt[2]{x} \log^2}$ for $x > 0$.
5. Show that for $x \geq 2$, $\pi(x) = \frac{\mathcal{J}(x)}{\log x} + \int_2^x \frac{\mathcal{J}(t)}{t \log^2 t} dt$.
6. Prove that $a \equiv b \pmod{m}$ if and only if a and b give the same remainder when divided by m .
7. Show that if a prime p does not divide a , then $a^{p-1} \equiv 1 \pmod{p}$.
8. Given that $m \geq 1$, $(a, m) = 1$ and $f = \exp_m(a)$. Show that $a^k \equiv a^h \pmod{m}$ if and only if $k \equiv h \pmod{f}$.

(5 × 1 = 5)

Turn over





G 18001031

Part B

*Answer any **five** questions.
Each question carries weight 2.*

9. For $n \geq 1$, prove that $\phi(n) = n \prod_{p|n} \left(1 - \frac{1}{p}\right)$.
10. State and prove the generalised inversion formula. Also deduce Generalised Mobius inversion formula.
11. Prove that the set of lattice points visible from the origin has density $6/\pi^2$.
12. Prove that the following relations are logically equivalent :

(a) $\lim_{x \rightarrow \infty} \frac{\pi(x) \log x}{x} = 1.$

(b) $\lim_{x \rightarrow \infty} \frac{\mathcal{J}(x)}{x} = 1.$

(c) $\lim_{r \rightarrow \infty} \frac{\Psi(x)}{x} = 1.$

13. Prove the following :

(a) $\hat{a} = \hat{b}$ if and only if $a \equiv b \pmod{m}$.

(b) Two integers x and y are in the same residue class if and only if $x \equiv y \pmod{m}$.

(c) The m residue classes $\hat{1}, \hat{2}, \dots, \hat{m}$ are disjoint and their union is the set of all integers.

14. Show that for any prime p , all the co-efficients of the polynomial

$$f(x) = (x-1)(x-2)\dots(x-p+1) - x^{p-1} + 1 \text{ are divisible by } p.$$

15. State and prove Chinese remainder theorem.





G 18001031

16. Let p be an odd prime and let d be any positive divisor of $p - 1$. Show that in every reduced residue system mod p , there are exactly $\phi(d)$ numbers a such that $\exp(a) = d$.

(5 × 2 = 10)

Part C

*Answer any **three** questions.
Each question carries weight 5.*

17. For all $x \geq 1$ show that :

$$\sum_{n \leq x} d(n) = x \log x + (2c - 1)x + O(\sqrt{x}), \text{ where } c \text{ is the Euler's constant.}$$

18. (a) Prove that for every $x \geq 1$, $[x]! = \prod_{p \leq x} p^{\alpha(p)}$ where the product is extended over all primes

$$\leq x \text{ and } \alpha(p) = \sum_{m=1}^{\infty} \left[\frac{x}{p^m} \right].$$

- (b) If $x \geq 2$ $\log [x]! = x \log x - x + O(\log x)$.

- (c) For $x \geq 2$ $\sum_{p \leq x} \left[\frac{x}{p} \right] \log p = x \log x + O(x)$.

where the sum is extended over all primes $\leq x$.

19. State and prove Shapiro's Tauberian theorem.

20. (a) State and prove Wolstenholme's theorem.

- (b) Show that the set of lattice points in the plane visible from the origin contains arbitrary large square gaps.

Turn over





G 18001031

21. (a) State and prove Lagrange's theorem for polynomial congruence.
- (b) Let f be a polynomial with integer co-efficients, let m_1, m_2, \dots, m_r be positive integers relatively prime in pairs, and let $m = m_1 m_2 \dots m_r$. Prove that the congruence $f(x) \equiv 0 \pmod{m}$ has a solution if and only if each of the congruences $f(x) \equiv 0 \pmod{m_i}$ ($i = 1, 2, \dots, r$) has a solution. Also show that if $V(m)$ and $V(m_i)$ denote the solutions of $f(x) \equiv 0 \pmod{m}$ and $f(x) \equiv 0 \pmod{m_i}$ for $i = 1, 2, \dots, r$, then $V(m) = V(m_1) V(m_2) \dots V(m_r)$.

22. If $|x| < 1$, prove that $\prod_{m=1}^{\infty} \frac{1}{1-x^m} = \sum_{n=0}^{\infty} p(n) x^n$, where $p(0) = 1$, $p(n)$ denotes the partition function.

(3 × 5 = 15)

